

Environmental influences on egg and clutch sizes in lentic- and lotic-breeding salamanders

Jon M. Davenport and Kyle Summers

Department of Biology and North Carolina Center for Biodiversity, East Carolina University, Greenville, NC 27858, USA.
E-mail: davenportjo05@students.ecu.edu.

Abstract

Environmental influences on egg and clutch sizes in lentic- and lotic-breeding salamanders. Recent research indicates that social and environmental factors influence egg and clutch sizes in amphibians. However, most of this work is based on the reproductively diverse order Anura (frogs and toads), whereas less research has been conducted on Caudata (salamanders) and Gymnophiona (caecilians). Researchers have suggested that a relationship exists between social and environmental factors and egg and clutch sizes in salamanders, but studies controlling for phylogenetic context are lacking. We could not identify a sufficient number of comparisons for social influences on egg and clutch sizes; therefore, we focused on environmental influences for this study. Data on egg size, clutch size, environmental factors, and phylogenies for salamanders were assembled from the scientific literature. We used independent, pair-wise comparisons to investigate the association of larval salamander habitat and egg size and the association of larval salamander habitat with clutch sizes within a phylogenetic framework. There is a significant association between larval habitat and egg size; specifically, stream-breeding species produce larger eggs. There is no significant association between larval habitat and clutch size. Our study confirms earlier reports that salamander egg size is associated with larval environments, but is the first to use phylogenetically independent contrasts to account for the lack of phylogenetic independence of the traits measured (egg size and clutch size) associated with many of the diverse lineages. Our study shows that environmental selection pressure can be quite strong on one aspect of salamander reproduction—egg size.

Keywords: Caudata, larval environment, independent contrasts, lentic, lotic, egg site deposition.

Received 5 April 2010.
Accepted 10 June 2010.
Distributed December 2010.

Resumo

Influência do ambiente sobre o tamanho do ovo e da desova em salamandras que se reproduzem em corpos d'água lênticos e lóticos. Pesquisas recentes indicam que fatores sociais e do ambiente influenciam o tamanho do ovo e da desova nos anfíbios. Contudo, a maioria desses trabalhos baseia-se na ordem Anura (anuros), que possui uma grande diversidade reprodutiva, e poucas pesquisas foram conduzidas nas ordens Caudata (salamandras) e Gymnophiona (cecílias). Sugeriu-se que existe uma relação entre fatores sociais e do ambiente e o tamanho do ovo e da desova em salamandras, mas não há estudos feitos dentro de um contexto filogenético. Não encontramos um número suficiente de comparações para a influência do comportamento social sobre o tamanho do ovo e da desova; portanto, neste estudo enfocamos a influência do ambiente. Os dados sobre tamanho do ovo e da desova, fatores do ambiente e filogenia das salamandras foram coletados na literatura. Usamos comparações *pair-wise* independentes na investigação da associação entre o habitat larval e o tamanho do ovo e entre o habitat larval e o tamanho da desova em um contexto filogenético. Encontramos uma associação significativa entre o habitat larval e o tamanho do ovo; especificamente, espécies que se reproduzem em riachos produzem ovos maiores. Não encontramos uma associação significativa entre o habitat larval e o tamanho da desova. Nosso estudo confirma relatos anteriores de que o tamanho do ovo das salamandras está relacionado ao ambiente em que as larvas se desenvolvem, mas é o primeiro a utilizar contrastes filogeneticamente independentes para explicar a ausência de independência filogenética das características medidas (tamanho do ovo e da desova) associadas a muitas das diferentes linhagens. Nosso estudo mostra que a pressão de seleção do ambiente pode ser muito intensa sobre um dos aspectos da reprodução das salamandras—o tamanho do ovo.

Keywords: Caudata, ambiente larval, contrastes independentes, ambientes lênticos, ambientes lóticos, local de ovipostura.

Introduction

Environmental factors and parental care play key roles in the evolution of offspring size and number (Shine 1978, Nussbaum and Shultz 1989, Clutton-Brock 1991, Stearns 1992). Shine (1978) proposed the “safe-harbor hypothesis,” which predicted that typically, the evolution of parental care should precede the evolution of larger eggs. Other researchers argued that environmental factors may influence the number and size of offspring more than parental care (Nussbaum 1985, Morrison and Hero 2003). Specifically, Nussbaum (1985, 1987) proposed that the evolution of larger eggs typically precedes the evolution of parental care.

Amphibians, particularly anurans, provide an excellent system with which to evaluate relationships among parental care, egg size, and environmental factors. Recent comparative

analyses of anurans confirmed a positive relationship between parental care and egg size (Summers *et al.* 2006, 2007). The authors found that egg size increased significantly with parental care. They also examined environmental factors and demonstrated that lotic- and montane-breeding anurans produce significantly larger eggs than their sister clades (Summers *et al.* 2007). These studies were the first to control for phylogenetic effects with regard to egg size in amphibian evolution.

Previous studies investigating relationships between environmental factors and egg and clutch sizes in salamanders lacked controls for phylogenetic effects (Salthe 1969, Kaplan and Salthe 1979). Coincidentally, Nussbaum's original proposal (1985, 1987) that the evolution of larger eggs should precede the evolution of parental care was based on salamander data. Nussbaum argued that larger eggs would be

avored in lotic environments (flowing water such as streams and seeps) because newly hatched larvae would have to consume larger food items in contrast to larvae hatched in lentic environments (standing water such as ponds or lakes). Larger eggs take longer to develop, enforcing a tradeoff between offspring size and number. Longer development would lead to increased mortality in the embryonic stage unless selection favored behaviors (e.g., parental care) to reduce egg mortality.

Comparative analyses require well-supported phylogenies, and well-supported salamander phylogenies have not been available until recently. We used multiple phylogenies (Methods below) to construct a phylogenetic supertree within which to evaluate relationships among egg size, clutch size, and environmental factors. Social influences on egg and clutch sizes could not be tested due to a lack of phylogenetic independence. Because most salamander parental care has evolved in one family (Plethodontidae), we could not make valid comparisons within those clades (Wells 2007). Therefore, we focus here on the hypothesis that environmental factors influence egg and clutch sizes in salamanders with larval development in lentic and lotic environments. Specifically, we concentrated on determining whether the site of egg deposition affects egg size and clutch size in this subset of salamanders. Based on the work by Nussbaum (1985), we predicted that egg size would be greater and clutch size would be smaller in taxa inhabiting lotic environments than in taxa inhabiting lentic environments.

Materials and Methods

We constructed a phylogenetic supertree consisting of all salamander species for which we could obtain both phylogenetic information and response-variable data to test our hypotheses within a phylogenetic framework. The supertree was constructed with the same hierarchical approach as Summers *et al.* (2007). We used a small number of studies that addressed specific phylogenetic relationships (e.g., relationships

within a genus). We preferentially chose studies that used DNA sequence data and maximum likelihood analyses. These studies were chosen because of (1) the quantity of sequence data available, (2) the thoroughness of the analyses, and (3) the well-found statistical rationale for the methods employed. The studies that were chosen consisted of the most recent studies of phylogenetic relationships of salamanders based on the same gene regions when possible.

The analysis utilized the following references for each group: Order Caudata (Wiens *et al.* 2005, Frost *et al.* 2006, Roelants *et al.* 2007); ambystomatids (Shaffer *et al.* 1991, Shaffer and Knight 1996, Weisrock *et al.* 2006a); salamandrids (Steinfartz *et al.* 2006, Weisrock *et al.* 2006b, Zhang *et al.* 2008); hynobiids (Zhang *et al.* 2006); plethodontids (Chippendale *et al.* 2004, Wiens *et al.* 2005).

Phylogenetic, pair-wise comparisons were used for our comparative analysis (Moller and Birkhead 1992, Maddison 2000, Summers *et al.* 2007). This method is used to compare continuous characters with discrete characters, and commonly is used in studies focusing on phylogenetic, pair-wise comparisons in which the traits of a focal group are compared to its sister clade (Gotmark 1994, Summers *et al.* 2007). Each phylogenetic pair-wise comparison was considered independent and counts of these points were analyzed with a χ^2 test with Yates correction when necessary.

Mean egg and clutch sizes were the continuous characters for this study. Egg size was defined as diameter of the ovum in millimeters after eggs were laid (including the jelly coat). The discrete characters in our analyses were larval salamander environments (lentic vs. lotic). Data on continuous characters (egg and clutch sizes) and discrete characters (larval environments) were taken from the primary literature for 31 species of salamanders (Appendices I and II); references from which data were extracted are listed in Appendix III. For species with multiple records for egg or clutch sizes, we averaged values (Appendix III). We also averaged egg and clutch

values for sister species inhabiting similar larval environments to facilitate comparison with closely related species that occupy different larval environments. Phylogenetic, pair-wise comparisons were conducted only on species for which we had all of the above data. Female body size was excluded from our analyses because no significant relationship was found between egg size and maximum female body size ($n = 25$, $R^2 = 0.0238$, $F = 0.56$, $p = 0.4619$). This was also true for egg size and mean female body size ($n = 25$, $R^2 = 0.0195$, $F = 0.46$, $p = 0.5058$).

Results

Eleven comparisons were identified for pair-wise comparisons (when controlling for phylogenetic effects) between lentic- versus lotic-breeding salamanders (Figure 1). We detected a significant association between the larval salamander environment and larger egg sizes ($\chi^2 = 9.00$, $df = 1$, $p < 0.05$; Table 1). Specifically, we found that egg sizes are significantly larger in lotic larval environments. We did not find a significant association between larval salamander environment and clutch size ($\chi^2 = 2.88$, $df = 1$, $p > 0.05$; Table 1).

Discussion

Our results partially agree with previous results relevant to the association between larval salamander environment and egg and clutch sizes. This association has been documented previously, but not in a phylogenetic context (Nussbaum 1985, 1987, 2003). Nussbaum (1985) originally hypothesized that lotic environments

would favor larger egg size, based primarily on prey items available for consumption by the larval salamanders; thus, lotic-breeding salamanders hatched at larger sizes to harvest the size classes of the most abundant food. This contrasts to the lentic-breeding salamanders with smaller embryos that hatch at smaller sizes and gorge on abundant, small zooplankton. Therefore the lotic environment lacking zooplankton would favor larger sizes of eggs and hatchlings. These results also are supported by recent work by Summers *et al.* (2007) with frogs in lotic environments that have larger egg sizes than those in lentic environments.

It is somewhat surprising that salamander clutch size was not found to be associated with the environment of larval salamanders in our study. This disparity may be the result of the small sample size for independent contrasts (9 comparisons). Nonetheless, salamanders that deposit eggs in lotic larval environments tend to have smaller clutch sizes than those that deposit eggs in lentic environments. This is in accord with previous work that hypothesized that adult female salamanders are subject to energy constraints. Thus, females that lay larger eggs deposit smaller clutches and females that lay smaller eggs deposit larger clutches (Nussbaum 2003). The variability of clutch sizes reported for some species may have contributed to the lack of association between larval environment and clutch size. Some species of salamanders in our study are widely distributed and exhibit geographic variation in egg and clutch sizes. For such species, we pooled data from across the range of the species, thereby increasing the variability of our data.

Table 1. Pair-wise comparisons of larval salamander deposition site and egg/clutch sizes.

	Sample Size	χ^2	<i>p</i> -value	Effect
Lotic versus Lentic – egg size	10 larger, 1 smaller	9.00	<0.05	Lotic larger
Lotic versus Lentic – clutch size	2 larger, 7 smaller	2.88	>0.05	—




Figure 1. Salamander phylogenetic supertree reconstructed from recently published phylogenies. Black branches are lentic-breeding species and white branches are lotic-breeding species included in our analysis. Salamander genus and species names can be found in Appendix I.


Parental care also may influence egg and clutch sizes; however, we could only identify three independent comparisons. In general, a lack of independent phylogenetic contrasts limited valid comparisons. In some families

(specifically Plethodontidae comprising 50% of extant salamanders), parental care is known for most species, making it difficult to compare social environments. Hence, for most of the taxa for which there are data on parental care, the

comparisons were not phylogenetically independent. Data on presence or absence of parental care are lacking for many species, especially the several new taxa that have been described recently (Camp *et al.* 2009). Detailed natural-history data are needed for many species, especially for the Asian and New World tropical salamanders, to test the parental care hypothesis properly.

Egg and clutch sizes were not associated with the same factors; thus, larger egg size has a significant association with lotic environments, but clutch size lacks a significant association with the environment defined. Because clutch size tends to be smaller in lotic-breeding species, salamander egg and clutch sizes may be responding to similar environmental selective pressures.

Acknowledgments

We thank Jason Brown, Michael Brewer and Jason Bond for assistance with constructing and editing the salamander phylogenetic supertree. We also thank Dave Beamer, Trip Lamb, and Linda Trueb for providing feedback on previous versions of this manuscript. JMD was supported by NSF Grant DEB-0716558. 

References

- AmphibiaWeb. 2010. Information on Amphibian Biology and Conservation. Berkeley, California: AmphibiaWeb. Electronic database accessible at <http://amphibiaweb.org/>. Captured on 05 April 2010.
- Anderson, P. L. 1943. The normal development of *Triturus pyrrhogaster*. *Anatomical Record* 86: 59–63.
- Anderson, J. D. and R. G. Webb. 1978. Life history aspects of the Mexican salamander *Ambystoma rosaceum* (Amphibia, Urodela, Ambystomatidae). *Journal of Herpetology* 21: 89–93.
- Anderson, J. D. and R. D. Worthington. 1971. The life history of the Mexican salamander *Ambystoma ordinarium* Taylor. *Herpetologica* 32: 165–176.
- Bishop, S. C. 1941. The Salamanders of New York. *New York State Museum Bulletin* 324: 1–365.
- Brame, A. H., Jr. 1956. The number of eggs laid by the California newt. *Herpetologica* 12: 325–326.
- Brame, A. H., Jr. 1968. The number of egg masses and eggs laid by the California newt, *Taricha torosa*. *Journal of Herpetology* 2: 169–170.
- Bruce, R. C. 1968. *Life History Studies of the Salamanders of the Genus Pseudotriton (Caudata: Plethodontidae)*. Unpublished Ph.D. Thesis. Duke University, Durham, NC.
- Bruce, R. C. 1969. Fecundity in primitive plethodontid salamanders. *Evolution* 23: 50–54.
- Bruce, R. C. 1971. Life cycle and population structure of the salamander *Stereochilus marginatus* in North Carolina. *Copeia* 1971: 234–246.
- Bruce, R. C. 1978a. Reproductive biology of the salamander *Pseudotriton ruber* in the southern Blue Ridge Mountains. *Copeia* 1978: 417–423.
- Bruce, R. C. 1978b. Life-history patterns of the salamander *Gyrinophilus porphyriticus* in the Cowee Mountains, North Carolina. *Herpetologica* 34: 53–64.
- Camp, C. D., W. E. Peterman, J. R. Milanovich, T. Lamb, J. C. Maerz, and D. B. Wake. 2009. A new genus and species of lungless salamander (family Plethodontidae) from the Appalachian highlands of the south-eastern United States. *Journal of Zoology* 279: 86–94.
- Chippendale, P. T., R. M. Bonnett, A. S. Baldwin, and J. J. Wiens. 2004. Phylogenetic evidence for a major reversal of life history evolution in plethodontid salamanders. *Evolution* 58: 2809–2822.
- Clergue-Gazeau, M. 1999. *Euproctus asper* (Duges 1852)—Pyrenäen-Gebirgsmolch. Pp. 251–269 in K. Grossenbacher and B. Thiesmeier (eds.), *Handbuch der Reptilien und Amphibien Europas. Band 4/1: Schwanzlurch (Urodela) I (Hynobidae, Proteidae, Plethodontidae, Salamandridae)*, AULA-Verlag, Wiebelsheim, Germany.
- Clutton-Brock, T. H. 1991. *The Evolution of Parental Care*. Princeton, NJ: Princeton University Press. 352 pp.
- Connon, F. E. 1947. A comparative study of the respiration of normal and hybrid *Triturus* embryos and larvae. *Journal of Experimental Zoology* 105: 1–24.
- Franz, R. 1964. The eggs of the long-tailed salamander from a Maryland cave. *Herpetologica* 20: 216.
- Frost, D. R., T. Grant, J. Faivovich, R. H. Bain, A. Haas, C. F. B. Haddad, R. O. de Sá, A. Channing, M. Wilkinson, S. C. Donnellan, C. J. Raworthy, J. A. Campbell, B. L. Blotto, P. Moler, R. C. Drewes, R. A. Nussbaum, J. D. Lynch, D. M. Green, and W. C. Wheeler. 2006. The Amphibian Tree of Life. *Bulletin of the American Museum of Natural History* 297: 1–370.

- Gallien, L. and M. Durocher. 1957. Table chronologique de development chez *Pleurodeles waltii* Michah. *Bulletin Biology France Belgique* 91: 97–114.
- Gibbons, J. W. and R. D. Semlitsch. 1991. *Guide to the Reptiles and Amphibians of the Savannah River Site*. University of Georgia Press, Athens, Georgia. 144 pp.
- Goin, C. J. 1951. Notes on the eggs and early larvae of three more Florida salamanders. *Annals of the Carnegie Museum* 32: 253–263.
- Goris, R. C. and N. Maeda. 2004. *Guide to the Amphibians and Reptiles of Japan*. Krieger Publishing Co., Malabar, FL. 285 pp.
- Gotmark, F. 1994. Are bright birds distasteful – a reanalysis of Cott, H.B. data on the edibility of birds. *Journal of Avian Biology* 25: 184–197.
- Haker, K. 1997. Haltung und Zucht des Chinesischen Riesensalamanders *Andrias davidianus*. *Salamandra* 33: 69–74.
- Hutchison, V. H. 1956. Notes on the plethodontid salamanders, *Eurycea lucifuga* (Rafinesque) and *Eurycea longicauda* (Green). *Occasional Papers of the National Speleological Society* 3: 1–24.
- Kaplan, R. H. and S. N. Salthe. 1979. The allometry of reproduction: an empirical view in salamanders. *American Naturalist* 113: 671–689.
- Licht, L. E. 1989. Reproductive parameters of unisexual *Ambystoma* on Pelee Island, Ontario. Pp. 209–217, in Dawley, R.M. and J.P. Bogart (eds.), *The Evolution and Ecology of Unisexual Vertebrates*. New York State Museum Bulletin, Number 466, Albany, New York.
- Liu, C. C. 1945. Life history of *Batrachuperus pinchonii*. *Journal of West China Border Research Society* 40: 44–55.
- Maddison, W. P. 2000. Testing character correlation using pairwise comparisons on a phylogeny. *Journal of Theoretical Biology* 202: 195–204.
- Minton, S. A., Jr. 1972. *Amphibians and Reptiles of Indiana*. *Indiana Academy of Sciences Monograph* 3: 1–346.
- Minton, S. A., Jr. 2001. *Amphibians and Reptiles of Indiana*. 2nd edition. Indiana Academy of Science, Indianapolis, Indiana. 404 pp.
- Moller, A. P. and T. R. Birkhead. 1992. A pairwise comparative method as illustrated by copulation frequency in birds. *American Naturalist* 139: 644–656.
- Morrison C. and J. M. Hero. 2003. Geographic variation in life-history characteristics of amphibians: a review. *Journal of Animal Ecology* 72: 270–279.
- Mosher, H. S., F. A. Fuhrman, H. D. Buchwald and H. G. Fischer. 1964. Tarichatoxin-tetrodotoxin: a potent neurotoxin. *Science* 144: 1100–1110.
- Noble, G. K. and J. A. Richards. 1932. *Experiments on the Egg-Laying of Salamanders*. American Museum Novitates, Number 513, American Museum of Natural History, New York.
- Nussbaum, R. 1985. The evolution of parental care in salamanders. *Miscellaneous Publications of the Museum of Zoology, University of Michigan* 169: 1–50.
- Nussbaum, R. 1987. The evolution of parental care in salamanders: an examination of the safe harbor hypothesis. *Researches on Population Ecology (Kyoto)* 29: 27–44.
- Nussbaum, R. 2003. Parental Care. Pp. 527–612 in B.G. M. Jamieson and D. Sever (eds.), *Reproductive Biology and Phylogeny of Urodela*. Science Publishers Inc., Enfield, NH, USA.
- Nussbaum, R. A. and D. L. Schultz. 1989. Coevolution of parental care and egg size. *American Naturalist* 133: 591–603.
- Petranka, J. W. 1998. *Salamanders of the United States and Canada*. Smithsonian Institution Press, Washington, D.C. 592 pp.
- Riemer, W. J. 1958. Variation and systematic relationships within the salamander genera *Taricha*. *University of California Publications in Zoology* 56: 301–390.
- Rocca, F. D., L. Vignoli, and M. A. Bologna. 2005. The reproductive biology of *Salamandrina terdigitata* (Caudata, Salamandridae). *Herpetological Journal* 15: 273–278.
- Roelants, K., D. J. Gower, M. Wilkinson, S. P. Loader, S. D. Biju, K. Guillaume, L. Moriau, and F. Bossuyt. 2007. Global patterns of diversification in the history of modern amphibians. *Proceedings of the National Academy of Sciences* 104: 887–892.
- Ryan, T. J. and R. C. Bruce. 2000. Life history evolution and adaptive radiation of hemidactyliine salamanders. Pp. 303–326 in Bruce, R. C., R. G. Jaeger and L. D. Houck (eds.), *The Biology of Plethodontid Salamanders*. Kluwer Academic/Plenum Publishers, New York.
- Salthe, S. N. 1969. Reproductive modes and the number and size of ova in the urodeles. *American Midland Naturalist* 81: 467–490.
- Schmidtler, J.J. and J.F. Schmidtler. 1975. Untersuchungen an westpersischen Bergbachmolchen der Gattung *Neurergus* (Caudata, Salamandridae). *Salamandra* 11: 84–98.

- Schwartz, A. and R. Etheridge. 1954. New and additional herpetological records from the North Carolina Coastal Plain. *Herpetologica* 10: 167–171.
- Semlitsch, R. D. and M. A. McMillian. 1980. Breeding migrations, population size structure, and reproduction of the dwarf salamander, *Eurycea quadridigitata*, in South Carolina. *Brimleyana* 3: 97–105.
- Shaffer, H. B., J. M. Clark, and F. Kraus. 1991. When molecules and morphology clash: A phylogenetic analysis of the North American ambystomatid salamanders (Caudata: Ambystomatidae). *Systematic Zoology* 40: 284–303.
- Shaffer, H. B. and M. L. McKnight. 1996. The polytopic species revisited: Genetic differentiation and molecular phylogenetics of the tiger salamander, *Ambystoma tigrinum* (Amphibia: Caudata) complex. *Evolution* 50: 417–433.
- Shine, R. 1978. Propagule size and parental care: the “safe harbor” hypothesis. *Journal of Theoretical Biology* 75: 417–424.
- Stearns, S. 1992. *The Evolution of Life Histories*. Oxford: Oxford Univ. Press. 264 pp.
- Steinfartz, S. 1995. Zur fortpflanzungsbiologie von *Neurergus crocatus* und *Neurergus strauchii barani*. *Salamandra* 31: 15–32.
- Steinfartz, S., S. Vicario, J. W. Arntzen, and A. Caccone. 2006. A Bayesian approach on molecules and behavior: reconsidering phylogenetic and evolutionary patterns of the Salamandridae with emphasis on *Triturus newts*. *Journal of Experimental Zoology Part B: Molecular and Developmental Evolution* 308: 139–162.
- Summers, K., C.S. McKeon, and H. Heying. 2006. The evolution of parental care and egg size: a comparative analysis in frogs. *Proceedings of the Royal Society of London B*. 273: 687–692.
- Summers, K., C.S. McKeon, H. Heying, J. Hall, and W. Patrick. 2007. Social and environmental influences on egg size evolution in frogs. *Journal of Zoology* 271: 225–232.
- Tago, K. 1931. *Newts and Salamanders*. Maruzen Co., Tokyo, Japan. 210 pp.
- Takahashi, H. and H. Iwasawa. 1990. Egg size variation among various breeding habitats in the salamander, *Hynobius lichenatus*. *Ecological Research* 5: 393–398.
- Tarkhishvili, D. N. and R. K. Gokhelasvili. 1999. The amphibians of the Caucasus. *Advances in Amphibian Research in the Former Soviet Union* 4: 1–239.
- Thiemeier, B. and C. Hornberg. 1998. Zur fortpflanzung von *Pachytriton labiatus*- ein weiterer Hinweis auf Brutpflegeverhalten in der Gattung *Pachytriton*. *Salamandra* 34: 77–80.
- Thorn, R. 1968. *Les Salamanders d'Europe, d'Asie, et d'Afrique du Nord*. Paul Lechevalier, Paris, France.
- Timofeev, B. I. 1997. Notes on the reproduction of *Neurergus crocatus* in captivity. *Advances in Amphibian Research in the Former Soviet Union* 2: 173–176.
- Twitty, V. C. 1935. Two new species of *Triturus* from California. *Copeia* 1935: 73–80.
- Twitty, V. C. 1936. Correlated genetic and embryological experiments on *Triturus* I and II. *Journal of Experimental Zoology* 74: 239–302.
- Weisrock D. W., H. B. Shaffer, B. L. Storz, S. R. Storz, and S. R. Voss. 2006a. Multiple nuclear gene sequences identify phylogenetic species boundaries in the rapidly radiating clade of Mexican ambystomatid salamanders. *Molecular Ecology* 15: 2489–2503.
- Weisrock D. W., T. J. Papenfuss, J. R. Macey, S. N. Litvinchuk, R. Polymeni, I. H. Ugurtas, E. Zhao, and A. Larson. 2006b. A molecular assessment of phylogenetic relationships and lineage accumulation rates within the family Salamandridae (Amphibia, Caudata). *Molecular Phylogenetics and Evolution* 41: 368–383.
- Wells, K. 2007. *The Ecology and Behavior of Amphibians*. University of Chicago, Chicago, USA. 1400 pp.
- Wiens, J. J., R. M. Bonett, and P. T. Chippendale. 2005. Ontogeny discombobulates phylogeny: Paedomorphosis and higher level salamander relationships. *Systematic Biology* 54: 91–110.
- Wood, J. T. and H. R. Rageot. 1963. The nesting of the many-lined salamander in the Dismal Swamp. *Virginia Journal of Science* 14: 121–125.
- Zhang, P., Y.-Q. Chen, H. Zhou, X.-L. Wang, T. J. Papenfuss, D. B. Wake, and L.-H. Qu. 2006. Phylogeny, evolution, and biogeography of Asiatic salamanders (Hynobiidae). *Proceedings of the National Academy of Sciences* 103: 7360–7365.
- Zhang, P., T. J. Papenfuss, M. H. Wake, L. Qu, and D. B. Wake. 2008. Phylogeny and biogeography of the family Salamandridae (Amphibia: Caudata) inferred from complete mitochondrial genomes. *Molecular Phylogenetics and Evolution* 49: 586–597.
- Zuffi, M. A. L. 1999. *Salamandrina terdigitata* (Lacépède 1788)—Brillensalamander. Pp. 29–246 in K. Grossenbacher and B. Thiesmeier (eds.), *Handbuch der Reptilien und Amphibien Europas. Band 4/1: Schwanzlurch (Urodela) I (Hynobiidae, Proteidae, Plethodontidae, Salamandridae)*, AULA-Verlag, Wiebelsheim, Germany.

Appendix I. Salamander species in our analyses and their larval environmental categories.

Salamander taxon	Larval environment	
Hynobiidae		
<i>Batrachuperus karlschmidti</i>	Lotic-stream	—
<i>Hynobius katoii</i>	Lotic-stream	—
<i>Hynobius lichenatus</i>	—	Lentic-pond
<i>Hynobius naevius</i>	Lotic-stream	Lentic-pond
<i>Salamandrella keyserlingii</i>		Lentic-pond
Salamandridae		
<i>Calotriton asper</i>	Lotic-stream	—
<i>Cynops ensacauda</i>	—	Lentic-pond
<i>Cynops pyrrhogaster</i>	—	Lentic-pond
<i>Neurergus crocatus</i>	Lotic-stream	—
<i>Neurergus kaiseri</i>	Lotic-stream	—
<i>Neurergus strauchii</i>	Lotic-stream	—
<i>Ommatriton vittatus</i>	—	Lentic-pond
<i>Pachytriton brevipes</i>	Lotic-stream	—
<i>Pachytriton labiatus</i>	Lotic-stream	—
<i>Pleurodeles walti</i>	—	Lentic-pond
<i>Salamandrina terdigitata</i>	Lotic-stream	—
<i>Taricha granulosa</i>		Lentic-pond
<i>Taricha rivularis</i>	Lotic-stream	—
<i>Taricha torosa</i>	—	Lentic-pond
<i>Triturus cristatus</i>	—	Lentic-pond
<i>Triturus marmoratus</i>	—	Lentic-pond
Plethodontidae		
<i>Eurycea longicauda</i>	Lotic-stream	—
<i>Pseudotriton ruber</i>	Lotic-stream	—
<i>Eurycea quadridigitata</i>	—	Lentic-pond
<i>Gyrinophilus porphyriticus</i>	Lotic-stream	—
<i>Pseudotriton ruber</i>	Lotic-stream	—
<i>Stereochilus marginatus</i>	—	Lentic-pond
Ambystomatidae		
<i>Ambystoma barbouri</i>	Lotic-stream	—
<i>Ambystoma rosaceum</i>	Lotic-stream	—
<i>Ambystoma ordinarium</i>	Lotic-stream	—
<i>Ambystoma texanum</i>	—	Lentic-pond
<i>Ambystoma tigrinum</i>	—	Lentic-pond

Appendix II. Salamander independent contrasts.

Lentic-pond	Lotic-stream
<i>Salamandrella keyserlingii</i>	<i>Batrachuperus karlschmidti</i>
<i>Ambystoma texanum</i>	<i>Ambystoma barbouri</i>
<i>Ambystoma tigrinum</i>	<i>Ambystoma ordinarium</i> , <i>A. rosaceum</i>
<i>Pleurodeles waltii</i>	<i>Salamandrina terdigitata</i>
<i>Taricha torosa</i> , <i>T. granulosa</i>	<i>Taricha rivularis</i>
<i>Cynops pyrrhogaster</i> , <i>C. ensacauda</i>	<i>Pachytriton brevipes</i> , <i>P. labiatus</i>
<i>Eurycea quadridigitata</i>	<i>Eurycea longicauda</i>
<i>Stereochilus marginatus</i>	<i>Pseudotriton ruber</i> , <i>Gyrinophilus porphyriticus</i>
<i>Hynobius lichenatus</i>	<i>Hynobius katoii</i> , <i>H. naevius</i>
<i>Ommatriton vittatus</i>	<i>Neurergus strauchii</i> , <i>N. kaiseri</i> , <i>N. crocatus</i>
<i>Triturus marmoratus</i> , <i>T. cristatus</i>	<i>Calotriton asper</i>

Appendix III. Salamander egg- and clutch-size data.

Taxon	Egg size	Clutch size	Authority
Cryptobranchidae			
<i>Andrias davidianus</i>	8.6	—	Haker 1997
Hynobiidae			
<i>Batrachuperus karlschmidtii</i>	3.7		Liu 1945
<i>Hynobius katoi</i>	4.8–5.0		AmphibiaWeb 2010
<i>Hynobius lichenatus</i>	2.8–3.2	40–60	Takahashi and Iwasawa 1990, Goris and Maeda 2004, AmphibiaWeb 2010
<i>Hynobius naevius</i>	5.0	34–72	Goris and Maeda 2004, AmphibiaWeb 2010
<i>Salamandrella keyserlingii</i>	1.5–2.0		Tago 1931
Salamandridae			
<i>Calotriton asper</i>	3.5–5.0	20–30	Clergue-Gazeau 1999, AmphibiaWeb 2010
<i>Cynops ensacauda</i>	2.8		Tago 1931
<i>Cynops pyrrhogaster</i>	2.0		Anderson 1943
<i>Neurergus crocatus</i>	1.5–2.0		Schmidtler and Schmidtler 1975, Timofeev 1997
<i>Neurergus kaiseri</i>	1.5–2.0		Schmidtler and Schmidtler 1975
<i>Neurergus strauchii</i>	2.6–3.0		Steinfartz 1995
<i>Ommatriton vittatus</i>	1.8–2.3		Tarkhnishvii and Gokhlashvii 1999
<i>Pachytriton brevipes</i>	3.5		Thorn 1968
<i>Pachytriton labiatus</i>	4.7–5.3		Thiemeier and Hornberg 1998
<i>Pleurodeles waltii</i>	1.7	150–1300	Gallien and Durocher 1957, AmphibiaWeb 2010
<i>Salamandrina terdigitata</i>	1.8	1–65	Zuffi 1999, Rocca <i>et al.</i> 2005
<i>Taricha granulosa</i>	1.8		Twitty 1936, Connon 1947
<i>Taricha rivularis</i>	2.8	10	Twitty 1935, 1936, Connon 1947, Riemer 1958
<i>Taricha torosa</i>	2.3	7–47	Twitty 1936, Connon 1947, Brame 1956, 1968, Mosher <i>et al.</i> 1964
<i>Triturus cristatus</i>	2.0	70–600	Thorn 1968, AmphibiaWeb 2010
<i>Triturus marmoratus</i>	2.0	200–380	Thorn 1968, AmphibiaWeb 2010

Appendix III. Continued.

Taxon	Egg size	Clutch size	Authority
Plethodontidae			
<i>Eurycea longicauda</i>	2.5–3.0	61–106	Hutchison 1956, Ryan and Bruce 2000, Minton 2001
<i>Eurycea quadridigitata</i>	1.8	7–62	Goin 1951, Semlitsch and McMillan 1980, Gibbons and Semlitsch 1991
<i>Gyrinophilus porphyriticus</i>	3.5–4.0	16–106	Bishop 1941, Bruce 1969, 1978b
<i>Pseudotriton ruber</i>	3.3	29–130	Bruce 1968, 1978a
<i>Stereochilus marginatus</i>	2.0–2.5	16–121	Richards 1932, Schwartz and Etheridge 1954, Wood and Rageot 1963, Ryan and Bruce 2000
Ambystomatidae			
<i>Ambystoma barbouri</i>	2.4–3.8	~260	Petranka 1998
<i>Ambystoma rosaceum</i>	2.6		Anderson and Webb 1978
<i>Ambystoma ordinarium</i>	2.8	109	Anderson and Worthington 1971
<i>Ambystoma texanum</i>	1.6–2.5	550–700	Minton 1972, Licht 1989, Petranka 1998
<i>Ambystoma tigrinum</i>	3.0	421–7000	Bishop 1941, Petranka 1998